



MICRO 90

FUNGICIDE - MITICIDE FOR SPRAYING

ACTIVE INGREDIENT:

Sulfur90.0%

INERT INGREDIENTS: 10.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with flowing water for at least 15 minutes. Get medical attention if irritation persists.

See panel below for additional "Precautionary Statements."

EPA Reg. No.55429-1

Manufactured by: **Continental Sulfur Company, LLC**

P.O Box 191 Kilgore, TX 75662

EPA Est. No. 55429-TX-1

Net Weight

Lbs. (__.__ kg)

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid inhalation of vapors, spray or dust. Avoid contact with eyes, skin and clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for wash ables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where run-off is likely to occur.

PHYSICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks or flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other workers, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI), of 24 hours. Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls, waterproof gloves,

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in its original labeled container in a cool, dry, locked place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TO USE AS A SPRAY

HOW TO MIX - Select a suitable hand sprayer or power unit which uses a ground rig or aerial applicator. Measure the recommended amount for the specific spray problem separately into water and stir to make a thin "slurry" paste. Fill the spray tank ½ full and then add the sulfur "slurry" to the tank before the other materials are added. When using a suitable "approved" wetting agent add it to the tank last to avoid possible foaming. Keep spray mixture agitated while spraying. Mix only the amount of spray needed for the job.

HOW MUCH SPRAY TO USE

WHEN TO APPLY THE LOWER RECOMMENDED RATE

1. During periods of minimal to moderate disease or insect infestation.
2. For regular season Cover Sprays.
3. When temperatures are up in the high 80s and low 90s.

WHEN TO APPLY THE HIGHER RECOMMENDED RATE

1. When mites or the diseases are damaging the crop and when weather conditions are favorable for their continued damage.
2. When temperatures are below and will not exceed 90°F within the following 3 days after the application.

AS A SPRAY - The lower rates would be 2 or 3 lbs. of the 2 to 6 lbs./100 gallons rate recommendation. The higher rates would be 4 or 6 lbs./100 gallons.

DIRECTIONS FOR DILUTION DILUTE APPLICATION

Ground: Apply specified rate in 20 to 60 gallons of water per acre.

Orchard: Apply specified rate in 100 to 800 gallons of water per acre.

CONCENTRATE APPLICATION

Ground: Apply specified rate in 5-10 gallons of water per acre.

Orchard: Apply specified rate in 20 to 100 gallons of water per acre.

AERIAL APPLICATION

Ground: Apply specified rate in 3 to 20 gallons of water per acre.

Orchard: Apply specified rate in 10 to 20 gallons of water per acre.

PRECAUTIONS

In order to obtain effective mite and disease control, both the upper and lower sides of plant foliage and fruit must be thoroughly covered with spray solution. Poor spray coverage will result in poor mite and disease control. When growing crops for "processing," consult the processor before applying sulfur. Do not use on any crop unless sulfur has been shown to be safe for that crop in your locality. Sulfur may burn foliage when temperatures exceed 90°F. Sulfur will cause severe fruit and leaf injury to sensitive plants. Use only on varieties known to be tolerant to sulfur. Do not apply within 14 days of oil spray for all crops except **CITRUS**. **FOR CITRUS** do not apply within 21 days of an oil spray.

AS A SPRAY FOR FRUIT AND NUTS

ALMONDS, PLUMS, PRUNES: For BROWN ROT, POWDERY MILDEW, LEAF SPOTS, SILVER MITES, FLAT MITES, SPIDER MITES. Use 10 to 30 lbs. per acre. Apply at bloom or early petal fall, repeat as necessary, usually 10 to 14 day intervals or after periods of wet weather for disease control. **FOR BROWN ROT** and **PRUNE ROT** on Plums and Prunes use 30 to 35 lbs. per acre 5 weeks before harvest. Repeat each 7 days if wet weather prevails and disease threatens during the harvest period.

APPLES AND PEARS: For POWDERY MILDEW, SCAB, BLISTER MITES, SILVER MITES & SPIDER MITES. Use 10 to 30 lbs. per acre. Apply Pre-Bloom and Petal Fall. Apply with cover sprays as needed throughout the season. For SCAB use 30 to 35 lbs. per acre at "Green Tip" and again at "Pink Bud." Pink Bud treatments will also control POWDERY MILDEW. Do not apply during or shortly before frost conditions. **PRECAUTION:** Certain varieties of apples and pears are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.

CITRUS: (GRAPEFRUIT ~ LEMONS, LIMES, ORANGES, TANGELOS, TANGEKINES, CITRUS CITRON, KUMQUATS, and hybrids of these). **For FLAT MITES, SILVER MITES, & SPIDER MITES: AS A SPRAY** use 5 to 10 pounds per 100 gallons of water for ground applications. Use 10 to 30 lbs. in 15 to 20 gallons of water per acre for aerial application. For Florida CITRUS RUST MITE use 25 lbs. per 500 gallons water equal to 40 to 50 lbs. sulfur per acre for ground applications. Use 40 to 50 lbs. in 15 to 20 gallons water per acre for aerial application. Otherwise, apply in any spray NOT CONTAINING OIL as needed for control during season or within 3 weeks of an oil spray as fruit burn may occur. Apply when mites are present in damaging numbers and repeat as needed.

PEACHES, NECTARINES: For BROWN ROT, POWDERY MILDEW, SILVER MITES, SPIDER MITES. Use 10 to 30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 days intervals or after periods of wet weather for disease control

. For POWDERY MILDEW use 50 pounds per acre at petal fall, 2 weeks later and again at beginning of pit hardening. For RUST and POWDERY MILDEW use 50 lbs. after pit hardening. For SILVER MITE Use 25 to 35 lbs. per acre after thinning. For BROWN ROT use 50 to 60 lbs. per acre 4 weeks pre-harvest and again 2 weeks later.

RASPBERRIES: For POWDERY MILDEW. Apply 6 to 15 lbs. per acre. Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires.

STRAWBERRIES: For POWDERY MILDEW and RED SPIDER MITES. Start in early leaf stage and repeat applications at 10 to 14 day intervals or as necessary for control with or without pesticide sprays. Use 3 to 20 lbs. per acre pre-harvest. See WHEN TO USE.

AS A SPRAY FOR VEGETABLES

BEANS, PEAS, BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE and PEPPERS for POWDERY MILDEW and on BEANS, PEAS AND PEPPERS for SPIDER MITES. Start in the early leaf stage. Repeat application every 10 to 14 days with or without pesticides or immediately after a rain. For POWDERY MILDEW & SPIDER MITES use 3 to 20 lbs. per acre. Do not apply to varieties susceptible to sulfur injury. See WHEN TO USE. MELONS: For POWDERY MILDEW use 25 lbs. per acre. For SPIDER MITES use 20 to 50 lbs. per acre. Do not apply to varieties susceptible to sulfur injury. Melon foliage is extremely sensitive to sulfur burn. Observe use precautions and apply only when temperatures are well below 90°F. See paragraphs on WHEN TO USE and HOW MUCH SPRAY OR DUST TO USE.

PEANUTS: AS A SPRAY for POWDERY MILDEW, SPIDER MITES and LEAF SPOT diseases apply 3 to 5 pounds per acre. Start in the early leaf stage and repeat applications as necessary or at 10 to 14 day intervals.

TOMATOES: For RUSSET MITES. Thorough coverage of upper and lower sides of leaves and fruit is required for effective control. Apply when mites are present in damaging numbers and repeat at 10 to 14 day intervals as needed. Use 30 to 40 lbs. per acre. If crop is being grown with the aid of stakes or poles, apply 10 to 13 lbs. per acre per foot of average plant height. (Plants 4ft. height = 40 to 52 lbs. pr acre.)

IMPORTANT

In order to obtain effective mite and disease control, both the upper and lower sides of plant foliage and fruit must be thoroughly covered with spray solution. Poor spray coverage will result in poor mite and disease control. When growing crops for "processing," consult the processor before applying sulfur. Do not use on any crop unless sulfur has been shown to be safe for that crop in your locality. Sulfur may burn foliage when temperatures exceed 90°F. Sulfur will cause severe fruit and leaf injury to sensitive plants. Do not use on or allow product to drift on to Apricots, D'Anjou or Comice Pears, Cranberries, Cucurbits (Cucumbers, Squash and some varieties of Melons), Filberts, Spinach, Tung Trees, Walnuts and other sensitive plants. Do not apply within 14 days of oil spray for all crops except citrus. **For Citrus do not apply within 21 days of an oil spray.**